X-ray Astronomy at SLAC¹

P. M. Saz Parkinson²

Stanford Linear Accelerator Center, Stanford University, Stanford, CA 94309

Abstract

The USA (*Unconventional Stellar Aspect*) experiment was launched in February of 1999 and operated for approximately 18 months. Group K at SLAC (Stanford Linear Accelerator Center) participated in this experiment along with the Naval Research Laboratory (NRL). I discuss the USA experiment and the data accumulated, along with some of the results obtained from the observations of such objects as the extragalactic BL Lac object 1ES1959+65, the Black Hole Candidate (BHC) XTE J1118+480, and the eccentric X-ray binary system Circinus X-1.

The USA experiment

USA is an X-ray timing experiment built jointly by the Naval Research Laboratory and the Stanford Linear Accelerator Center. USA was launched on 1999 February 23 on the Advanced Research and Global Observation Satellite (ARGOS), into a nearly circular 830 km Sun-synchronous orbit at 98.8° inclination.

USA is sensitive in the energy range 1–15 keV with an effective area of about $1000~\rm cm^2$ at 3 keV and a field of view which is approximately 1.2° FWHM circular. The Crab Nebula gives about $3600~\rm cts\,s^{-1}$ at the center of the field of view. Most observations were taken in an event mode, with 32 μ s time resolution and 16 pulse height channels. For more details on the USA instrument see [Ray et al.(1999)].

USA Observations

Table 1 shows the total amount of USA data accumulated. The data is grouped by object class. For each class I list the top three sources observed by USA.

1ES1959+65

USA observed this BL Lac object, at the time a potential TeV emitter, and detected variability on the timescale of a few days with a three-fold flux increase, consistent with what has been observed in the optical band at other times. A correlation was observed between the X-ray flux and the spectrum, with the spectrum hardening as the source becomes brighter [Giebels et al.(2002)].

¹Work at SLAC was supported by Department of Energy contract DE-AC 03-76-SFO0515. Basic research in X-ray astronomy at the Naval Research Laboratory is supported by ONR/NRL. This work was also supported by the NASA Applied Information Systems Research Program

² for the USA collaboration. e-mail: pablos@SLAC.stanford.edu

maintaining the data needed, and coincluding suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding and DMB control number.	tion of information. Send commen narters Services, Directorate for In:	ts regarding this burden estimate formation Operations and Reports	or any other aspect of the s, 1215 Jefferson Davis	nis collection of information, Highway, Suite 1204, Arlington	
1. REPORT DATE DEC 2002		2. REPORT TYPE		3. DATES COVE 00-00-2002	RED 2 to 00-00-2002	
4. TITLE AND SUBTITLE			5a. CONTRACT NUMBER			
X-ray Astronomy at SLAC				5b. GRANT NUMBER		
				5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
					5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Stanford University, Stanford Linear Accelerator Center, Stanford, CA, 94309					8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITO	RING AGENCY NAME(S) A	10. SPONSOR/MONITOR'S ACRONYM(S)				
					11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAIL Approved for publ		ion unlimited				
13. SUPPLEMENTARY NO	TES					
14. ABSTRACT						
15. SUBJECT TERMS						
16. SECURITY CLASSIFIC	ATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 2	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE				

Report Documentation Page

Form Approved OMB No. 0704-0188

Table 1: Summary of USA observations

	· J
Exposure (ks) a	Examples b
655	Mkn 421 (300), 3C273 (120), 1ES1959+650 (60)
1835	Cyg X-1 (700), XTE J1118 (320), GRS 1915 (290)
765	X0142+614 (250), $E2259+586$ (205), $1E1048.1$ (120)
1255	Crab (600), PSR 1509-58 (205), PSR B0540-69 (90)
2680	Cyg X-2 (355), Cir X-1 (235), EXO 0748 (230)
1300	SMC X-1 (315), Cen X-3 (210), LMC X-4 (170)
	655 1835 765 1255 2680

^aapproximate amount of raw data accumulated

XTE J1118+480

The black hole transient XTEJ1118+480 was discovered by RXTE during the second of two outbursts that it underwent in 2000 [Remillard et al.(2000)]. USA observed the source from 10 April to 11 June of 2000 and tracked the evolution of a low frequency (~0.1 Hz) QPO, showing that the QPO frequency is not correlated with the source flux [Wood et al.(2000)].

Circinus X-1

Circinus X-1 is currently classified as a NS due to the detection of Type I X-ray bursts by EXOSAT in the 1980s. We studied the timing and spectral evolution of Cir X-1 around its 16.6 day orbit. We modeled its energy spectrum with two components, one representing the disk and one representing a Comptonizing region. We studied the numerous QPO present in Cir X-1 and determined that their origin probably lies in the Comptonizing region, rather than the disk. We compared the power spectrum of Cir X-1 to that of the BHC Cyg X-1 and found that if scaled by 3.8, the two breaks in both spectra coincide. If these breaks are inversely proportional to the mass, this would imply a mass for Cir X-1 in excess of the canonical mass for a neutron star [Saz Parkinson (2003)].

References

[Giebels et al. (2002)] Giebels, B. et al. 2002, ApJ, 571, 763

[Ray et al.(1999)] Ray, P. S. et al. 1999, Proceedings of X-ray Astronomy 1999, Bologna, Italy, AIP Conference Proceedings, 599, 336 (2001), astro-ph, 9911236

[Remillard et al.(2000)] Remillard, R., Morgan, E., Smith, D., & Smith, E. 2000, IAU CIRC, 7389, 2

[Saz Parkinson (2003)] Saz Parkinson, P. M. 2003, PhD Thesis, Stanford University

[Wood et al.(2000)] Wood, K. S. et al. 2000, ApJL, 544, L45

^bTop three sources with the most amount of data, shown in ks in parentheses